












Oratect® Quick Reference Guide

The Oratect® Oral Fluid Drug Screen Device is a simple one-step test for the detection of drugs of abuse in oral fluid.

<p>1. Remove the blue cap by holding the sides and pulling gently. This will expose the collection pad. Make sure there is a blue line present in each window area indicating an unused device.</p> 	<p>2. Open mouth and gently rub the collection pad inside mouth against cheek in a circular motion several (approximately 15-20) times. Make sure to keep head level.</p> 	<p>3. Gently rub the collection pad against the opposite cheek in circular motion several (approximately 15-20) times.</p> 
<p>4. Gently rub the collection pad on top of the tongue several (approximately 15-20) times. Do not chew, suck, bite or bend the collection pad.</p>  <p>5. Rub the collection pad underneath the tongue several (approximately 15-20) times.</p>	<p>6. Place the collection pad underneath the tongue for approximately 30 seconds to collect saliva. Instruct the donor to hold the device in place with hand. When sufficient amount of saliva is collected the blue lines will move. Repeat steps 2-6 until blue lines start to move upward</p> 	<p>7. Remove from mouth as soon as blue lines move in both of the test windows. Re-cap the device.</p>  <p>8. Lay the device on a flat surface and read results in approximately 5 minutes after removing the device from mouth. Read package insert.</p>

Interpreting Test Results

<p>Invalid Result When no colored band appears in the CONTROL (C) region, the test is invalid even if there is a band in the test region. Repeat the test with a new device.</p>  <p>Example Interpretation:</p> <p>ME: Invalid TH: Invalid CO: Invalid AM: Invalid OP: Invalid PC: Invalid</p>	<p>Negative Result For each test, two colored bands should be observed:</p> <ul style="list-style-type: none"> • One in the CONTROL (C) region • One in the specific TEST region <p>The color of the test band may be slightly darker or lighter than the control band. Any visible band that can be seen is a negative result.</p>  <p>Example Interpretation:</p> <p>ME: Negative TH: Negative CO: Negative AM: Negative OP: Negative PC: Negative</p>	<p>Presumptive Positive Result A colored band at the CONTROL (C) region should be observed. When there is no colored band or shadow band at the specific TEST region, the test is presumptive positive for that particular drug.</p>  <p>Example Interpretation:</p> <p>ME: Presumptive Positive TH: Negative CO: Negative AM: Negative OP: Negative PC: Negative</p>
--	---	--